


# PL-TBD Resin



## Description

Polymer supported 1,5,7-triazabicyclo[4.4.0]dec-5-ene

## Synonyms



1,3,4,6,7,8-Hexahydro-2*H*-pyrimido[1,2-*a*]pyrimidine resin

## Applications

PL-TBD Resin is a polymer supported strong base, which can be employed where MTBD would otherwise be employed. It can be used to promote or accelerate reactions, remove excess reagents or remove acidic by-products.

Care should be taken to ensure that the free base form and the resultant salt complex are both well solvated in the reaction solvent.

PL-TBD MP is a macroporous material suitable for use with protic, polar and apolar solvents. This reagent can also be used to effectively scavenge Platinum, Palladium and Tin species from organic solution.



## References

- Tomoi, M et al (1984), Makromol Chem, **185**, 2117
- Tomoi, M et al (1985), Makromol Chem, **186**, 2473
- Iijima, K et al (1992), Pure Appl Chem, **A29**, 249
- Schuchardt, U et al (1996), J Mol Catal A Chem, **109**, 37
- Caldarelli, M et al (1999), J Chem Soc, Perkin Trans 1, 107
- Boisnard, S et al (1999), Tetrahedron Lett, **40**, 7469
- Organ, M G & Dixon, C E (2000), Biotechnol & Bioeng (Comb Chem), **71**, 71
- Simoni, D et al (2000), Tetrahedron Lett, **41**, 1607
- Alhambra, C et al (2001), Tetrahedron Lett, **42**, 6675

## Products Information

Microporous

### PL-TBD Resin

2.7mmol/g 150-300µm (50-100 mesh)

Macroporous

### PL-TBD MP-Resin

1.7mmol/g 100Å 150-300µm (50-100 mesh)